Total number of printed pages-7

3 (Sem-2/CBCS) PHY HC 2

## 2023

## PHYSICS

(Honours Core)

(Waves and Optics)

Paper: PHY-HC-2026

Full Marks: 60

Time: Three hours

The figures in the margin indicate full marks for the questions.

- 1. Answer the following questions:  $1 \times 7 = 7$ 
  - (a) What is the relation between group velocity  $v_g$  and wave velocity v in a dispersive medium?
  - (b) What is the nature of wavefront emitted by a point source?

- (c) Which method is used for producing two coherent sources from one single source in Newton's rings experiment?
  - (d) What is the grating element for a plane diffraction grating having 5,00,000 lines/cm?
  - (e) What do you mean by a positive zone plate?
  - (f) What is the velocity of a particle at the nodes of a standing wave?
  - (g) Which assumption was considered by Newton while formulating the velocity of sound as incorrect?
- 2. Answer the following questions: 2×4=8
  - (a) Fundamental frequency of a stretched string of length 50 cm and mass 10 gm is 300 Hz. What is the tension applied?

- (b) What are the conditions essential to obtain sustained interference of light?
- (c) In Fraunhofer diffraction pattern formed by a single slit, suppose that the slit width is 0.03 cm and the wavelength of light used is  $6 \times 10^{-5}$  cm. Find the diffraction angle for the first dark band.
- (d) Show that two perpendicular SHMs of equal frequency and equal amplitude but having a phase difference of  $\pi/2$  can produce a circular motion.
- 3. Answer any three of the following questions: 5×3=15
  - (a) Deduce an expression for the velocity of transverse vibrations in a stretched string.

- (b) Explain the phenomenon of refraction of a plane wave at a plane surface using Huygens' principle.
- (c) Illustrate Stokes treatment for explanation of the change of phase when reflection takes place at the denser medium.
  - (d) Mention three differences between Fresnel and Fraunhofer diffraction. A zone plate behaves like a convex lens of focal length 50 cm. If the wavelength of light is 5000 Å, calculate the radius of first half period zone.

    3+2=5
- (e) What do you mean by standing (stationary) waves? Deduce an equation illustrating the relationship between phase and group velocities. 1+4=5

- 4. Answer **any three** of the following questions:
  - (a) Determine the resultant of two perpendicular SHMs having frequency ratio 2:1 and a phase difference zero.

    Obtain a representation of the resultant path graphically.

    6+4=10
  - (b) Discuss the phenomenon of Fraunhofer diffraction at a single slit. Find an expression for the width of the central maximum. Fraunhofer diffraction pattern due to a narrow slit of width 0.2 mm is observed in a screen placed on the focal plane of a lens having focal length 2 m. If the first minima is at 5 mm on either side of central maximum, calculate the wavelength of the incident light. 7+3=10

(c) Describe Fresnel's biprism experiment for interference. How can you determine the wavelength of light by this method? Light of wavelength 5896 Å falls normally on a thin wedge-shaped air film forming fringes that are 3 mm apart. Find the angle of the wedge.

2+5+3=10

- (d) Find the expression for intensity due to a plane diffraction grating. Why cannot the secondary maxima be observed? What is its resolving power?

  5+2+3=10
- (e) Elucidate the construction and working principle of a Michelson's interferometer.

  Under what conditions are circular fringes formed in Michelson's interferometer? How are localized fringes formed in Micheleson's interferometer?

6+2+2=10

- (f) Write short notes on **any two** of the following:
  - (i) Ripple and gravity waves
  - (ii) Vibrations in a plucked string
  - (iii) Haidinger fringes
  - (iv) Holography